

FIRST RESPONDERS HOW TO DON PPE USING N95 RESPIRATOR

First Responder:		CDC-recommended PPE
Trained Observer:	Date:	1. 2. 6
PURPOSE: This checklist has been developed to assi Personal Protective Equipment (PPE), and securing or	ist First Responders in the proper donning and doffing of n-scene used PPE.	TWO PAIRS OF SINGLE-USE COVERALLS INVITH OR WITHOUT INTEGRATED SOCIO
HOW TO USE: This document should be used by Trained Observers and First Responders in conjunction with: The instructional First Responder Ebola Briefing + PPE PowerPoint published November 13, 2014. CDC training videos for donning and doffing PPE (www.cdc.gov/vhf/ebola/hcp/ppe-training/). CDC Interim Guidance for Emergency Medical Services (EMS) Systems and 9-1-1 Public Safety Answering Points (PSAPs) for Management of Patients with Known or Suspected Ebola Virus Disease in the United Sates – October 28, 2014. (www.cdc.gov/vhf/ebola/hcp/interim-guidance-emergency-medical-services-systems-911-public-safety-answering-points-management-patients-known-suspected-united-states.html). CDC Guidance on Personal Protective Equipment To Be Used by Healthcare Workers During Management of Patients with Ebola Virus Disease in U.S. Hospitals, Including Procedures for Putting On (Donning) and Removing (Doffing) – October 20, 2014. (www.cdc.gov/vhf/ebola/hcp/procedures-for-ppe.html). IMPORTANT NOTES:		(DISPOSABLE) NITRILE EXAM GLOVES (At a minimum, outer gloves should have extended cuffs) AND APRON AND BOOT or SHOE COVERS (SHOE COVERS ACCEPTABLE ONLY IF COVERALLS HAVE BYTEGARDIS SOCIO) (All should be single-use (disposable) fluid vesistant or imperior AND APRON AND BOOT or SHOE COVERS (All should be single-use (disposable) fluid vesistant or imperior AND APRON AND BOOT or SHOE COVERS (All should be single-use (disposable) fluid vesistant or imperior AND APRON AND BOOT or SHOE COVERS (All should be single-use (disposable) fluid vesistant or imperior AND APRON AND BOOT or SHOE COVERS (All should be single-use (disposable) fluid vesistant or imperior AND APRON AND BOOT or SHOE COVERS (SHOE COVERS ACCEPTABLE ONLY IN COVERALLS HAVE AND APRON AND BOOT or SHOE COVERS AND
Donning and doffing procedures are different, depending on the type of PPF selected		EBOLA PREPAREDNESS DEPARTMENT OF HEALTH AND HOSPITALS
Prior to donning, designate a Work Zone for doffing. Work Zone should allow a clear separation between clean and potentially contaminated areas and should be located as close to the Hot Zone as safely feasible .		
Step 1. Engage Trained Observer	and supervision of a Trained Observer.	
Step 2. Remove Personal Clothing and Items	☐ In a clean area, change into surgical scrubs (or footwear with closed toe and heels, and no hole Remove personal items (e.g., jewelry, watches, cell ph	
Step 3. Inspect PPE Prior to Donning	 ✓ Visually inspect: PPE ensemble to ensure it is in serviceable condition. All required PPE and supplies are available. Trained Observer reviews the donning sequence in a step-by-step fashion. 	Sizes selected are correct. be before beginning the donning process, reading aloud.
Step 4. Perform Hand Hygiene	Perform hand hygiene with alcohol-based hand	rub (ABHR). Allow hands to dry before next step.
Step 5. Put on Inner Gloves	Put on first pair of gloves. Cuffs should extend	as far up the arms as possible.
Step 6. Put on Boot or Shoe Covers	Boot covers should extend to mid-calf. Shoe covers are acceptable ONLY if used with	a coverall with integrated socks .
Step 7. Put on Coverall	Ensure coverall: — Is large enough for unrestricted freedom of movement. If tape is needed to secure sleeves over inner g	Cuffs of inner gloves are tucked under the sleeve of the coverall.
Step 8. Put on N95 Respirator	□ For a secure fit ensure: □ Top tie or elastic band is over the head and above the ear: □ Bottom tie or elastic band goes along the back of the necl □ Complete a user seal check. Adjust mask so th	under the ears.
Step 9. Put on Hood	Hood goes over the N95 respirator. Hood MUST fully cover all hair, neck and ears and extend past the neck to the shoulders.	
Step 10. Put on Outer Apron (if used)	Put on full-body outer apron to provide additional protection to the front of the body. Apron should extend to mid-calf .	
Step 11. Put on Outer Gloves	Put on second pair of gloves (with extended cuffs). Ensure cuffs are pulled over coverall sleeves, extending as far up arms as possible. Using different color gloves when double gloving makes it easier to see contamination, cuts or tears.	
Step 12. Put on Face Shield	Ensure face shield is a full face shield ; place of This process provides additional protection to the shield is a full face shield ; place of This process provides additional protection to the shield is a full face shield; place of the shield is a full face shield; place of the shield is a full face shield; place of the shield is a full face shield; place of the shield is a full face shield; place of the shield is a full face shield; place of the shield is a full face shield; place of the shield is a full face shield; place of the shield is a full face shield; place of the shield is a full face shield; place of the shield is a full face shield; place of the shield is a full face shield; place of the shield is a full face shield; place of the shield is a full face shield; place of the shield is a full face shield in the sh	ver the N95 respirator AND the hood . he front and sides of the face , skin and eyes .
	☐ Trained Observer verifies:	

Donning process is **complete** and the **integrity** of the ensemble.

range of movement while all areas of the body remain covered.

☐ Disinfect outer-gloved hands with ABHR. Allow to dry prior to patient contact.

First Responder is comfortable and able to extend arms, bend at the waist, and go through a range of motions to ensure sufficient

Step 13. Verify

Step 14. Disinfect Outer Gloves